

**PLACE
PICTURE
HERE**

Name: _____ D.O.B.: _____

Allergy to: _____

Weight: _____ lbs. Asthma: **Yes (higher risk for a severe reaction)** **No**

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____

THEREFORE:

If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.

If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING:

SEVERE SYMPTOMS

 LUNG Shortness of breath, wheezing, repetitive cough	 HEART Pale or bluish skin, faintness, weak pulse, dizziness	 THROAT Tight or hoarse throat, trouble breathing or swallowing	 MOUTH Significant swelling of the tongue or lips
 SKIN Many hives over body, widespread redness	 GUT Repetitive vomiting, severe diarrhea	 OTHER Feeling something bad is about to happen, anxiety, confusion	<p>OR A COMBINATION of symptoms from different body areas.</p>

↓ ↓ ↓

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS

 NOSE Itchy or runny nose, sneezing	 MOUTH Itchy mouth	 SKIN A few hives, mild itch	 GUT Mild nausea or discomfort
--	--	--	--

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM


Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____


HOW TO USE AUVI-Q® (EPINEPRHINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case.
2. Pull off red safety guard.
3. Place black end of Auvi-Q against the middle of the outer thigh.
4. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
5. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN® AND EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
6. Remove and massage the injection area for 10 seconds.
7. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALICK®), USP AUTO-INJECTOR, IMPAX LABORATORIES

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip.
3. Grasp the auto-injector in your fist with the red tip pointing downward.
4. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh.
5. Press down hard and hold firmly against the thigh for approximately 10 seconds.
6. Remove and massage the area for 10 seconds.
7. Call 911 and get emergency medical help right away.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____

PHONE: _____

NAME/RELATIONSHIP: _____

PHONE: _____



Family Food Allergy Health History Form

Student Name: _____ Date of Birth: _____
Parent/Guardian: _____ Today's Date: _____
Home Phone: _____ Work: _____ Cell: _____
Primary Healthcare Provider: _____ Phone: _____
Allergist: _____ Phone: _____

1. Does your child have a diagnosis of an allergy from a healthcare provider: [] No [] Yes

2. History and Current Status

a. What is your child allergic to? [] Peanuts [] Insect Stings [] Eggs [] Fish/Shellfish [] Milk [] Chemicals [] Latex [] Vapors [] Soy [] Tree Nuts (walnuts, pecans, etc.) [] Other:
b. Age of student when allergy first discovered:
c. How many times has student had a reaction? [] Never [] Once [] More than once, explain:
d. Explain their past reaction(s):
e. Symptoms:
f. Are the food allergy reactions: [] Same [] Better [] Worse

3. Trigger and Symptoms

a. What are the early signs and symptoms of your student's allergic reaction? (Be specific; include things the student might say.)
b. How does your child communicate his/her symptoms?
c. How quickly do symptoms appear after exposure to food(s)? ____secs. ____mins. ____hrs. ____days
d. Please check the symptoms that your child has experienced in the past:
Skin: [] Hives [] Itching [] Rash [] Flushing [] Swelling (face, arms, hands, legs)
Mouth: [] Itching [] Swelling (lips, tongue, mouth)
Abdominal: [] Nausea [] Cramps [] Vomiting [] Diarrhea
Throat: [] Itching [] Tightness [] Hoarseness [] Cough
Lungs: [] Shortness of breath [] Repetitive Cough [] Wheezing
Heart: [] Weak pulse [] Loss of consciousness

4. Treatment

a. How have past reactions been treated?
b. How effective was the student's response to treatment?
c. Was there an emergency room visit? [] No [] Yes, explain:
d. Was the student admitted to the hospital? [] No [] Yes, explain:
e. What treatment or medication has your healthcare provider recommended for use in an allergic reaction?
f. Has your healthcare provider provided you with a prescription for medication? [] No [] Yes
g. Have you used the treatment or medication? [] No [] Yes
h. Please describe any side effects or problems your child had in using the suggested treatment:

5. Self Care

a. Is your student able to monitor and prevent their own exposures?	<input type="checkbox"/> No	<input type="checkbox"/> Yes
b. Does your student:		
1. Know what foods to avoid	<input type="checkbox"/> No	<input type="checkbox"/> Yes
2. Ask about food ingredients	<input type="checkbox"/> No	<input type="checkbox"/> Yes
3. Read and understands food labels	<input type="checkbox"/> No	<input type="checkbox"/> Yes
4. Tell an adult immediately after an exposure	<input type="checkbox"/> No	<input type="checkbox"/> Yes
5. Wear a medical alert bracelet, necklace, watchband	<input type="checkbox"/> No	<input type="checkbox"/> Yes
6. Tell peers and adults about the allergy	<input type="checkbox"/> No	<input type="checkbox"/> Yes
7. Firmly refuses a problem food	<input type="checkbox"/> No	<input type="checkbox"/> Yes
c. Does your child know how to use emergency medication?	<input type="checkbox"/> No	<input type="checkbox"/> Yes _____
d. Has your child ever administered their own emergency medication?	<input type="checkbox"/> No	<input type="checkbox"/> Yes _____

6. Family / Home

a. How do you feel that the whole family is coping with your student's food allergy?	_____
b. Does your child carry epinephrine in the event of a reaction?	<input type="checkbox"/> No <input type="checkbox"/> Yes
c. Has your child ever needed to administer that epinephrine?	<input type="checkbox"/> No <input type="checkbox"/> Yes
d. Do you feel that your child needs assistance in coping with his/her food allergy?	_____

7. General Health

a. How is your child's general health other than having a food allergy?	_____
b. Does your child have other health conditions?	_____
c. Hospitalizations?	_____
d. Does your child have a history of asthma?	<input type="checkbox"/> No <input type="checkbox"/> Yes
If yes, does he/she have an Asthma Action Plan?	<input type="checkbox"/> No <input type="checkbox"/> Yes
e. Please add anything else you would like the school to know about your child's health:	_____ _____

8. Notes:

Parent / Guardian Signature: _____ Date: _____

Reviewed by R.N.: _____ Date: _____